

## Solution to Exercise 3.2-3

Illustration

How large will be the distance  $d$  between the (center of gravity) of the positive and negative charges for reasonable field strengths and atomic numbers, e.g. the combinations of

- 1 kV/cm
- 100 kV/cm
- 10 MV/cm
- , the last one being about the ultimate limit for the best dielectrics there are,
- $z = 1$  (H, Hydrogen)
- $z = 50$  (Sn (= tin), ...)
- $z = 100$  (?)

From the backbone we have a relation for  $d$  as a function of  $z$ ,  $m$  the radius  $R$  of the atom, and the field strength  $E$ :

$$d_E = \frac{4 \pi \epsilon_0 \cdot R^3 \cdot E}{ze}$$

We need to look up some number for the radius of the three atoms given (try this link), then the calculation is straight forward - let's make a table:

Atom	$R$	$d(1 \text{ kV/cm})$	$d(100 \text{ kV/cm})$	$d(10 \text{ MV/cm})$
$z = 1$				
$z = 50$				
$z = 100$				

Compared to the radius of the atoms, the separation distance is tiny. No wonder, electronic polarization is a small effect *with spherical atoms!*

Calculate the "spring constant" and from that the resonance frequency of the "electron cloud" (assume the nucleus to be fixed in space).

If you don't know off-hand the resonance frequency of a simple harmonic oscillator - that's fine. If you don't know exactly what that is, and where you can look it up - you are in deep trouble.

Anyway, in [this link](#) you get all you need. In particular the resonance (circle) frequency  $\omega_0$  of a harmonic oscillator with the mass  $m$  and the spring constant  $k_S$  is given by

$$\omega_0 = \left( \frac{k_S}{m} \right)^{1/2}$$

How large are the spring constants? That is question already answered in the backbone, so we import the equation

$$k_S = \left( \frac{(ze)^2}{4 \pi \epsilon_0 \cdot R^3} \right)$$

Again, let's make a table for the answers:

Atom	Spring constant	$\omega_0$
<b>z = 1</b>		
<b>z = 50</b>		
<b>z = 100</b>		